Check for updates

Deep-ocean polymetallic nodules as a resource for critical materials

James R. Hein¹¹¹², Andrea Koschinsky² and Thomas Kuhn³

Abstract | Deep-ocean polymetallic nodules form on or just below the vast, sediment-covered, abyssal plains of the global ocean. Polymetallic nodules primarily consist of precipitated iron oxyhydroxides and manganese oxides, onto which metals such as nickel, cobalt, copper, titanium and rare earth elements sorb. The enormous tonnage of nodules on the seabed, and the immense quantities of critical metals that they contain, have made them a target for future mining operations. Mining of polymetallic nodules has been spurred by the need for critical metals to support growing populations, urbanization, high-technology applications and the development of a green-energy economy. Nevertheless, an improved understanding of the affected ecosystems and their connectivity, as well as the environmental impacts of deep-ocean mining, is required before operations begin. Opportunities exist, however, to ensure that this new industry applies adaptive management to continually refine operations with the goal of environmental protection and invests in the development of green technologies for extractive metallurgy and mining. In this Review, we explore the chemical processes that control the concentration of critical metals in deep-ocean polymetallic nodules, discuss the mining and metallurgical techniques required, and highlight the opportunities and potential risks that are presented by this new industry.

Deep-ocean polymetallic nodules (also known as manganese nodules) are composed of iron and manganese oxides that accrete around a nucleus on the vast abyssal plains of the global ocean¹⁻⁶. Polymetallic nodules vary in diameter from less than one to tens of centimetres and acquire economically interesting quantities of critical metals (metals that are essential to the security and economic wellbeing of a nation) from ocean water and/or sediment pore waters. As a result, the enormous quantities of these nodules - which are conservatively estimated to total 21 billion dry tons in the Clarion-Clipperton Zone7,8 (CCZ; located in the northeast equatorial Pacific Ocean) alone (FIG. 1) — host considerable tonnages of critical metals that are essential for green energy, vehicles and infrastructure, as well as for new technology and military applications. This reservoir of critical metals has, therefore, triggered interest in new deep-ocean mining operations. However, the potential environmental impacts of such activities are of great concern and have hastened extensive research and evaluation in this area^{9–11}.

Polymetallic nodules were first discovered on the seabed at a depth of approximately 4,300–5,500 m during the 1872–1876 voyage of the HMS Challenger to study the deep ocean¹². A number of nodules, which contained numerous shark teeth and cetacean ear bones, were collected from the western end of the CCZ nodule field¹². Early hypotheses suggested that polymetallic nodules form by alteration of volcanic rocks and grains, especially glass, which are transformed into manganese carbonates and, finally, into oxides that are deposited from solution to form nodules on water-rich sediments^{12–14}. However, later work — aided in part by the 1957–1958 International Geophysical Year — revealed the general source of the metals (seawater and sediment pore water) and the sorption processes that might be involved in the acquisition of minor metals by nodules^{15–20}.

Economic interest in the deep-ocean polymetallic nodules developed in the 1960s^{15,16}, which subsequently led to the formation of consortia in Germany, the United States, Canada, Japan, France, Belgium, Italy and the Netherlands in the 1960s and 1970s²¹⁻²³. These consortia aimed to develop resource assessments and extraction technologies for polymetallic nodules in the CCZ²¹. Deep-ocean-mining initiatives were developed to the point of several pilot mining operations^{21,24}, although most commercial activities ceased by the early 1980s, owing to low metal prices on the global market. Nevertheless, scientific interest in polymetallic nodules continued and geological, oceanographic, environmental and geochemical baseline studies were undertaken4,25-29. The Deep Ocean Mining Environmental Study programme, for example, involved 12 National Oceanic and Atmospheric Administration

¹U.S. Geological Survey, Pacific Coastal and Marine Science Center, Santa Cruz, CA, USA.

²Department of Physics & Earth Sciences, Jacobs University Bremen, Bremen, Germany.

³Federal Institute for Geosciences and Natural Resources (BGR), Hannover, Germany.

➡e-mail: jhein@usgs.gov
https://doi.org/10.1038/
s43017-020-0027-0

Key points

- Polymetallic nodules cover vast areas of the abyssal ocean floor and contain significant amounts of critical metals.
- The chemical and mineralogical compositions of polymetallic nodules are primarily controlled by their formation process.
- A unique characteristic of deep-ocean nodules compared to terrestrial deposits is the presence of multiple commodities in one deposit; for example, nodules from the Clarion–Clipperton Zone contain Mn, Ni, Cu and Co.
- Deep-ocean mining might avoid some of the environmental issues associated with terrestrial mining.
- The development of societies towards a more sustainable future cannot proceed without critical metals. Deep-ocean mining can not only deliver the metals necessary for this transition but can do so with a low carbon footprint.
- The precautionary approach, adaptive management and best environmental practices are essential to the development of a polymetallic nodule resource.

research cruises during 1975 and 1976, which integrated data from multiple disciplines to understand the distribution, formation and evolution of polymetallic nodules in three specific areas representative of the vast CCZ. The Deep Ocean Mining Environmental Study programme revealed that hiatuses in sedimentation, moderate to low primary productivity in surface water, Antarctic bottom-water flow and many other environmental conditions favour the formation of metal-rich nodules²⁵.

Over the following four decades, research continued throughout the central Pacific and provided new insights into, for example, the complex surfacechemical processes that allow for the acquisition of remarkably large metal contents and the local, regional and global environmental controls on the formation of alternating, micrometre-scale, Fe-rich and Mn-rich laminae^{1,2,10,24,30-43} (FIG. 2).

Research activities were controlled by exploration contracts with the International Seabed Authority, which was created based on the United Nations Convention on the Law of the Sea as an independent agency to administer mineral resources in areas beyond national jurisdictions (ABNJ; that is, all areas oceanward of the exclusive economic zones — 200 nautical miles from the coastline of coastal or island nations). Originally, research was predominantly undertaken by national agencies, until 2010, when private companies became involved and a polymetallic-nodule-mining industry was born. These exploration activities are ongoing today and polymetallic-nodule mining is likely to begin in earnest in the mid-to-late 2020s.

In this Review, we outline the chemical properties and formation characteristics of polymetallic nodules, before evaluating their economic significance and potential for future deep-ocean-mining. We also highlight the environmental and ecological significance of this developing industry and offer an overview of the opportunities and challenges that still remain. A comprehensive review of deep-ocean polymetallic nodules and their resource potential is critical, as these nodules provide an important natural laboratory for understanding chemical sedimentology and associated geochemical processes. In addition, deep-ocean polymetallic nodules represent a resource for critical elements (including Ni, Cu, Mn, Co, Mo, Ti, Li and rare earth elements, plus yttrium, or REY), which require secure and diverse sources to aid transition from a petroleum-based-energy world to a low-carbon future.

Nodule formation and coverage

Deep-ocean polymetallic nodules, composed of alternating Fe-rich and Mn-rich laminae, are predominantly formed on the sediment-covered floor of the global ocean, at water depths of approximately 3,500–6,500 m (FIG. 2a). In the following sections, we review the principle formation mechanisms of deep-ocean polymetallic nodules and outline how their chemistry and mineralogy (and, therefore, ore grade) define different types of potential ore deposits.

Nodule characteristics and formation. Deep-ocean polymetallic nodules have been subdivided based on their size (small, <4 cm diameter; medium, 4–8 cm; and large, >8 cm)^{1,30,45} and seafloor density (from tens up to a thousand nodules per square metre; FIG. 2a displays a region of the central CCZ covered by medium-sized nodules with a nodule density of ~20 kg m⁻²)¹. Polymetallic nodules form on, or just below, the sediment surface of the abyssal ocean by two principle processes, hydrogenetic and diagenetic precipitation^{4,46,47}. Both processes require a nucleus (for example, a shark's tooth) around which hydrogenetic and diagenetic layers can precipitate^{6,28,30,47} (FIG. 3).

Hydrogenetic precipitation is driven by the oxidation of dissolved Mn^{2+} and Fe^{2+} in oxygen-rich ocean waters and subsequent accretion of Mn^{4+} and Fe^{3+} oxide colloids around a nucleus^{32,47,48}. X-ray diffraction reveals that these precipitates are composed of amorphous ferrihydrite (FeOOH) and cryptocrystalline Fe-vernadite (δ -MnO₂), which possess positive and negative surface charges, respectively^{1,32,48}. Hydrogenetic layers form at very slow rates of a few millimetres per million years, which requires negligible sedimentation, as well as oxic conditions, on the ocean floor and in the underlying sediment^{1,4,7,24,30,34,39}.

Diagenetic precipitation, on the other hand, occurs within the pore space of deep-ocean sediments^{4,46}. Oxidation of organic matter in deep-ocean sediments results in the reduction and dissolution of Mn oxides and release of associated elements (Ni, Cu, Li, among others)^{1,4,36}. Owing to concentration gradients in the sediment, these metals diffuse upwards and, on contact with oxygen-rich ocean water, are reoxidized, leading to the precipitation of 7-Å and 10-Å Mn oxides (disordered phyllomanganates)^{1,2,4,36}. Diagenetic layers, which grow at a few tens of millimetres per million years, require slightly higher sedimentation rates (<10 mm kyr⁻¹) than hydrogenetic layers, reflecting climatically induced variations in surface-water biological productivity, as well as suboxic conditions on the ocean floor and in the underlying sediments1,4,7,46,47.

Note, phyllomanganates are sometimes referred to in the literature as buserite and birnessite^{49,50}, but those names correctly refer to ordered rather than disordered phyllomanganates. Mn oxides in deep-ocean





polymetallic nodules are made up of topotactic 7-Å and 10-Å manganates (which should strictly be termed vernadites)⁵¹. Topotactic means that the MnO₆ octahedral layers, which are stacked in the crystallographic *c* direction, are slightly offset to each other in the crystallographic *ab* plane. The minerals buserite and birnessite do not show this characteristic, as discussed elsewhere¹.

How polymetallic nodules remain on the sediment surface when their growth rates are so low compared to the surrounding sediments (3–5 mm kyr⁻¹) is still an unsolved paradox^{52–54}. Benthic biological activity and/or seismic shaking may represent key processes^{52–54}. Nevertheless, erosion and hiatuses in sedimentation are recognized as critical processes in the onset of nodule growth^{55–57}, whereas increased sedimentation rates and/or deposition of ash layers⁵⁷ may, ultimately, lead to the burial of a nodule generation.

It is important to note that ocean-floor nodules only represent the youngest generation of polymetallic nodules; hydroacoustic data reveal the presence of older nodule generations below the sediment surface^{58,59}. Older nodule generations might be a common phenomenon beneath nodule fields worldwide but require relatively oxygenated conditions (free oxygen in the range of a few tens of μ moll⁻¹) in the sedimentary column to prevent dissolution^{1,40}.

Nodules found on and below the ocean floor represent an important but unexploited resource of a wide range of critical metals. Before considering the processes by which these nodules can be collected, and the metals of interest extracted, it is important to consider how the formation processes described above influence their chemistry and, therefore, metal grade.

Formation controls on nodule chemistry. The Mn oxides and Fe oxyhydroxides that form the framework of the deep-ocean polymetallic nodules are able to efficiently scavenge a large number of dissolved elements from ocean water, which are only present in ultra-trace concentrations. As a result, polymetallic nodules typically possess high concentrations of minor and trace elements (for example, Li, Co, REY).

Recent synchrotron-based methods, such as X-ray absorption near-edge structure and extended X-ray absorption fine structure spectroscopy, built on prior statistical analysis, experimental work and thermodynamic models to demonstrate that metal enrichment and the structural incorporation of minor and trace metals in deep-ocean polymetallic nodules is controlled by the mineralogy of the nodule⁶⁰⁻⁶⁵. As a result, noduleformation processes - hydrogenetic versus diagenetic precipitation — that control the mineralogy of the polymetallic nodules also govern their chemical composition^{1,2,39,41}. As outlined above, very different geological and oceanographic conditions are required for these two processes. Therefore, variations in sedimentation rates and the extent of metal cycling through suboxic sediment resulting from factors such as increased microbiological activity represent the base-level controls on nodule chemistry (FIG. 3).

The Fe oxyhydroxide and Mn oxide colloids formed by hydrogenetic precipitation possess slightly positive



Fig. 2 | Small-scale distribution and structure of polymetallic nodules. a | Plan view of the seafloor in the central Clarion–Clipperton Zone at 4,900 m water depth with even coverage of about 20 kg m⁻² (dry weight) of medium-sized nodules. b | X-ray scan of a nodule cross section from the Clarion–Clipperton Zone with typical alternation of Mn-rich and Fe-rich layers. Sediment within the nodules is an indication of their high porosity. Image in part a courtesy of T. Kuhn, Federal Institute for Geosciences and Natural Resources (BGR), Germany.

and negative surface charges, respectively. The surface charge of these colloids, under ocean-water conditions, results in attraction, surface adsorption and enrichment of dissolved critical metals on the nodule surfaces^{32,41,42,66}. Cations (such as Cu²⁺, Ni²⁺ and Co²⁺) are attracted by the negatively charged δ -MnO₂ and anions (such as $UO_2(CO_3)_2^{2-}$) by the slightly positively charged FeOOH (FIG. 3a). In addition, the low charge density of the FeOOH surface enables the formation of covalent bonds with other slightly charged or neutral dissolved species, including Ti(OH)₄⁰ (REFS^{32,42,43}). Typically, hydrogenetic precipitates contain similar amounts of Fe and Mn (Fe/Mn ratio ~1 in bulk analyses), low Y/Ho and high Th/U ratios, and very high concentrations of Co, Te, Ce and Pt (REFS^{8,20,41}). The high concentrations of Co, Te, Ce and Pt result from oxidation of these elements on the surface of δ -MnO₂ particles, which represents a very efficient process for enriching trace metals that can overcome the electrostatic repulsion of negatively charged species in solution (for example, for Te and Pt) by the negatively charged Mn-oxide surface^{32,42,66-68}.

On the other hand, the 7-Å and 10-Å disordered phyllomanganates that characterize diagenetic nodules typically incorporate metal ions such as Ni²⁺, Cu²⁺ and Li⁺ to balance negative charge deficits in their crystal lattice^{1,2,33,34,51,66,68,69}. Charge deficits form as a result of structural defects (gaps) or the incorporation of Mn²⁺ and Mn³⁺ instead of Mn⁴⁺ (REFS^{1,34,46,63,68}). Diagenetic layers of polymetallic nodules are characterized by high Mn/Fe ratios (>10 in bulk analyses), high Ni, Cu and Li concentrations, low Co and Ce concentrations, high Y/Ho ratios and low Th/U ratios^{1,33,34} (FIG. 3). Metals of

REVIEWS

economic interest (Co, Ni, Cu, Mo, Zn) either replace Mn or occupy vacant sites in the octahedral MnO_6 layers of the Mn oxides^{1,33,34} (FIG. 3a). Interlayer sites of the 7-Å and 10-Å phyllomanganates are, by contrast, characterized by the presence of water molecules as well as alkali (such as Li, Na, K) and alkaline earth (such as Mg, Ca, Sr) elements^{1,33,34,70}.

In addition, X-ray absorption near-edge structure and extended X-ray absorption fine structure measurements further indicate that redox-sensitive elements, such as Co, Ce and Te, which are found bound in hydrogenetic nodules and crusts, are present in a higher oxidation state than in the overlying ocean water. This observation confirms that the exceptionally high concentration of such metals in polymetallic nodules results from redox processes involving Mn oxide^{31,35,60,64,65,68}. By contrast, other elements (such as Pb) remain divalent during surface adsorption and incorporation⁶⁸. Transition-metal cations and some anionic metal species form stable inner-sphere complexes on the surface of Mn and Fe oxides, which can lead to their isotopic fractionation. For example, Mo isotopes are fractionated during adsorption as molybdate, due to variations in the coordination environment of dissolved (tetrahedral coordination as MnO²⁻₄) and adsorbed (distorted octahedral coordination) Mo (REF.⁶²). Other species, for example, alkaline earth metals and anions such as chromate and selenate, form predominantly outer-sphere complexes on oxide surfaces, which are not associated with isotopic fractionation⁶¹. Hence, this molecular-scale information forms the basis for understanding the mechanisms by which critical metals are enriched in nodules to levels that foster economic interest.

The mean chemical compositions of polymetallic nodules in the best-known nodule fields can be used to define three types of potential ores^{8,56,66,71} (TABLE 1). The CCZ and Central Indian Ocean Basin nodules, which represent mixed hydrogenetic-diagenetic nodules, have similar compositions, with high Mn, very high Ni and Cu, and moderate Co, Mo and Li contents^{8,56,66}; nodules from the Peru Basin, which primarily form through diagenetic processes, are characterized by high Mn and very high Ni and Li contents^{8,66}; and the Penrhyn Basin nodules, which primarily form through hydrogenetic precipitation, have the highest Ti, Co, Y and total rare earth elements content of the three nodule types⁷¹ (TABLE 1). Therefore, these different nodule fields represent potential ores for a range of different critical metals, with the exact composition of the nodules (and, as a result, the metals that are of economic importance) controlled by the formation processes of each nodule field. Other elements with a high concentration relative to the Earth's crust are listed in TABLE 1 and may be available for extraction in the future.

Nodule distribution and exploration contract areas.

The most well-studied nodule fields include the CCZ, Penrhyn-Samoa Basin, Peru Basin, Central Indian Ocean Basin and abyssal plains surrounding seamounts and ridges in the west Pacific^{1,2,7,8,66} (FIG. 1). Although most nodule fields are in the ABNJ, important nodule deposits that represent large metal resources can also be found within the national jurisdictions of several Pacific Island countries and coastal nations, including Japan, Cook Islands, Kiribati, Niue and the United States^{57,71–73} (FIG. 1).</sup> Permissive areas for nodule formation are defined as those with sedimentation rates <10 mm per 1,000 years, a source of material for nucleation of nodules (such as particles of rock debris, indurated sediment, nodule fragments and shark teeth), surface waters that have moderate to low productivity and small-scale oceanfloor topography^{1,8,39,47,66} (FIG. 1). Permissive areas do not necessarily contain economically minable nodule deposits but simply represent regions where their formation is feasible.

Most contracts for exploration of marine mineral resources in the ABNJ are focused on the extraction of



Fig. 3 | Formation mechanisms and environments for polymetallic nodules.

a | Schematic showing the formation of δ -MnO₂ and ferrihydrite (bottom) and phyllomanganates (top), which precipitate around a nucleus to form hydrogenetic and diagenetic nodule layers, respectively. In addition to those shown here, many other anions and neutral complexes sorb onto the ferrihydrite and cations sorb onto the δ -MnO₂ and other phyllomanganates. **b** | Schematic showing the formation of hydrogenetic, mixedtype and diagenetic nodules, with variations in chemistry shown. Figure adapted with permission from REFS^{36,128}, Mineralogical Society of America. polymetallic nodules. At present, 18 contracts exist for nodule exploration, each covering 75,000 km², of which 16 are focused on nodule exploration in the northeast Pacific CCZ. Owing to the remarkable ability of polymetallic nodules to scavenge critical metals from ocean water, they represent a key resource of in-demand metals. As such, economic interest in deep-sea polymetallic nodules has boomed, necessitating deeper investigation into their exploration, mining and environmental impacts.

Mining deep-ocean mineral deposits

The European Union recognized that raw materials are crucial to Europe's economy and sustainable development and, as of 2017, identified a list of 27 critical raw materials (CRM). In 2018, the United States Geological Survey produced an updated list of 35 CRM essential to the economic and national security of the United States. Identifying reliable and sustainable sources of these CRM represents one of the primary factors driving the development of the deep-ocean-mining industry, especially as many CRM are at high risk of supply disruption, owing to the lack of diversity of their worldwide production⁷⁴. Typically, each CRM comes from just a few countries. For example, 60% of the current global production of raw Co, which is an indispensable component in lithiumion batteries, comes from the Democratic Republic of the Congo and the dominant supplier of refined Co is China (with an 80% market share)75. Increased diversity of CRM production and refining operations will be essential to ensure a secure supply of CRM for lowcarbon applications, such as green energy and green vehicles. Nevertheless, at the start of this new industry, it is important to consider the environmental and ecological impacts of deep-ocean mining, including how these can be minimized moving forward^{8,36,73,76}. In this section, we focus on the practical side of polymetallic-nodule mining (in terms of its rationale, incentives, challenges, environmental implications and technologies).

Unique (or favourable) characteristics of deep-ocean

mining. One of the first factors to be considered, with respect to potential deep-ocean mining, is whether the unique characteristics of marine-based mine sites provide favourable incentives for mine development. Marine-based mine sites do not require roads, oceanfloor ore-transport systems, water-transportation or electrical-transport systems, buildings, waste dumps or other infrastructure that characterize terrestrial mines. In addition, no overburden will need to be removed before mining can take place, as the deposits of interest are exposed at the ocean floor. Other crucial drivers of deepsea mining include the fact that many of the deposits that are present at a single marine mine site contain three or more metals of economic interest^{8,10,11}. Small deposits can be selectively mined simply by moving the productionmining vessel from one high-grade deposit to another, without the need to process intervening low-grade material. Therefore, compared to terrestrial mines, less ore is required to provide the same amount of metal^{8,10,11}. Acid mine drainage and river or soil contamination will be avoided by deep-ocean mining, as will many other challenges faced by terrestrial mine sites, such as the

Table 1 Mean composition of polymetallic nodules in four nodul	e fields
--	----------

Element (units)	CCZ	CIOB	Peru Basin	Cook Islands– Penrhyn Basin
lron/manganese ratio	0.22	0.21	0.25	0.96
Manganese (wt%)	28.4	24.4	34.2	16.9
lron (wt%)	6.16	7.14	6.12	16.20
Nickel (wt%)	1.30	1.10	1.30	0.377
Copper (wt%)	1.070	1.040	0.599	0.231
Titanium (wt%)	0.320	0.420	0.160	1.280
Cobalt (wt%)	0.210	0.111	0.048	0.375
Molybdenum (ppm)	590	600	547	295
TREE (ppm)	717	931	334	1537
Vanadium (ppm)	445	497	431	504
Zirconium (ppm)	307	752	325	555
Thallium (ppm)	199	347	129	146
Lithium (ppm)	131	110	311	51
Yttrium (ppm)	96	108	69	141
Arsenic (ppm)	67	150	65	150
Tungsten (ppm)	62	92	75	59

Data obtained from: CCZ^{8,56,66}, CIOB^{8,66}, Peru Basin^{8,66}, Cook Islands–Penrhyn Basin⁷⁰, CCZ, Clarion–Clipperton Zone; CIOB, Central Indian Ocean Basin; TREE, total rare earth elements.

relocation of towns and villages, deforestation and largescale lowering of the groundwater table. Deep-ocean mining also offers the prospect of reduced risk to on-site workers and the absence of child labour. Many of these issues are pervasive challenges in land-based mining, particularly in developing countries^{8,10,11}.

Comparisons with land-based deposits. In addition to the advantages presented by the unique characteristics of deep-ocean-mine sites, it is important to consider how the tonnage of critical metals hosted within polymetallic nodules compares to that found within terrestrial ore deposits. The mean metal content of nodules (TABLE 1), and the estimated tonnage of nodules within the entire CCZ, can be used to calculate the tonnage of each metal hosted in deep-ocean polymetallic nodules compared to land-based resources (summarized in TABLE 2). CCZ nodules are shown to contain a greater tonnage of Mn, Ni, Co, Tl and Y (and a similar tonnage of As) than the entire global terrestrial reserve base. The nodule tonnage of Tl, for example, is approximately 6,000 times higher than the global terrestrial reserve base and 6.5 times higher than the global terrestrial resource (TABLE 2). TABLE 2 also highlights metals that represent terrestrial-dominant resources (especially Cu, Ti and total rare earth elements) and those that represent marine-dominant resources (especially Co, Ni, Tl and Te).

Enormous quantities of metal evidently exist on the ocean floor in the CCZ, as well as other nodule fields. However, not all of that metal is economically extractable. Even for decades ahead, only a very small part of that enormous tonnage of nodules will be extracted. Nevertheless, TABLE 2 demonstrates that deepocean polymetallic nodules represent important resources for critical metals, which may be extracted in the future. However, before the mining and extraction of nodule-bound critical metals can occur, exploration is needed to identify economically viable, first-generation nodule-mine sites.

Exploration techniques. Exploration carried out by different groups since ~1980 showed that two main parameters can be used to distinguish between economic and uneconomic deposits: nodule abundance in kilograms per square metre (tonnage; cut-off ~10 dry kg m⁻² in the CCZ) and metal content in weight percent (cut-off, which will vary depending on tonnage and global markets, is ~2 wt.% Ni+Cu+Co for the CCZ)^{3,5,7,71}.

Polymetallic nodule exploration therefore requires, as a first step, the detection and delineation of large nodule fields in water depths between 3,500 and 6,500 m. Detection of nodule fields is generally achieved using multibeam echo-sounding systems mounted on the hull of research vessels^{3,77,78} (FIG. 4). In addition to bathymetric information, such systems provide acoustic imagery (backscatter) that can be used to interpret the geological conditions on the ocean floor, including the separation of flat, sediment-covered areas that contain nodules from areas devoid of them^{45,79-84}. After areas of interest (which can cover several thousand square kilometres) have been identified, a detailed bathymetric survey is conducted using autonomous underwater vehicles or deep-towed acoustic systems, such as side-scan sonar. Underwater systems survey at tens of metres up to about 100 m above the ocean floor, thus providing higher resolution (less than a few metres) bathymetric surveys than achievable from vessel-mounted systems (50–100 m)^{82,83}. High-resolution acoustic survey are typically accompanied by video mapping using either deep-towed video or photo sledges, remotely operated vehicles or autonomous underwater vehicles3 (FIG. 4). Sample collection, which is essential, is achieved using box corers (FIG. 4). This tool consists of a ~0.13-m³ metal box that samples a block of undisturbed sediment from the seafloor with polymetallic nodules still in place¹. As box corers sample a defined seafloor area, the mass of nodules per unit area can be measured directly. Moreover, representative samples are used to determine the nodule grade and metal content using optical emission spectrometry, mass spectrometry and other analytical techniques^{33,34}.

Data gathered by sample collection and highresolution acoustic surveys are then fed into multivariate (geo)statistical models to predict nodule abundance and metal resources over large areas^{5,78,79,83,84}. Such predictive models are based on artificial neural networks or random forests, as well as on classical geostatistics, such as variography and kriging, in order to assess the quantity of the resource^{5,83,85–87}. Once the economic potential of a nodule field has been determined, then, if favourable, the operations will move on to mining.

Mining techniques. Polymetallic-nodule-mining operations in the deep ocean consist of the following major components: collector robot on the ocean floor, vertical riser system, surface mining vessel, bulk carrier vessels at sea and processing plant on land^{1,11} (FIG. 4).

Consortia from South Korea, India, China and Europe have been developing several of these components up to a semi-industrial level⁸⁸⁻⁹².

Since polymetallic nodules lie either on the ocean floor or within the upper 20 cm of deep-ocean sediments, they can be harvested in a manner similar to potatoes in the field. Thus, mining machines might collect nodules from the ocean floor by either a mechanical or hydraulic uptake mechanism^{1,91,93,94}. The mechanical machinerv uses fork-like tools mounted on drums, which rotate backwards. By contrast, hydraulic uptake uses a water jet to lift nodules, together with ambient sediments, which are subsequently sucked into the nodule collector. Within the collector, sediment is cleaned from the nodules, which are then crushed to a certain size and pumped to a buffer chamber at the lower end of the vertical riser system^{90,91,95}. This buffer chamber ensures the continuous flow of nodules into the riser pipe. Vertical transport of nodules is achieved using either centrifugal pumps or air lift^{93,95}. At present, there is a major focus on the development of these pumping systems to provide improved flow-rate control and prevention of plugging from density waves⁹⁵. The full-scale vertical riser system must have a capacity of 300-500 tons of ore per hour and include critical links to the collector system on the ocean floor and the surface mining vessel⁹⁵. Collector locomotion on the ocean floor is achieved using a caterpillar track system in most mining machines. The surface mining vessel provides power to, and control over, all subsea systems, as well as housing ore-receiving, storage and ship-to-ship transfer capacities. Bulk carrier vessels with

at least 50,000 tons storage capacity are required for the shipment of nodules to shore⁹⁶.

Extractive metallurgy. Following the mining and collection of polymetallic nodules from the abyssal plain of the deep ocean, as well as transport to onshore facilities, the metals hosted in these nodules will need to be extracted. Development and upscaling of metallurgical routes for the processing of polymetallic nodules are essential prerequisites for the commercial viability of nodule mining. To ascertain the commercial viability and to increase public acceptance, the metallurgical processing of these nodules must lead to zero or minimal waste, a low carbon footprint, recycling of reagents and additives, and capital and operating costs that are similar to existing commercial operations. Because polymetallic nodules are a special type of oxide deposit that do not possess a terrestrial analogue, the development of new and green methods for metallurgical processing of polymetallic nodules is necessary and has been the subject of considerable research during the past few decades (discussed elsewhere97).

Since nodules are built up of nanometre-scale manganese oxides and iron oxyhydroxides that are epitaxially intergrown (crystallographically aligned FeOOH and MnO₂ laminae), it is not possible to separate and enrich the metals of interest from the other nodule components by conventional beneficiation methods, such as flotation, density separation or magnetic separation^{97,98}. The polymetallic-nodule matrix has to be completely destroyed to release the metals. This destruction can be

Table 2 Global tonnage of metals in CCZ polymetallic nodules versus terrestrial deposits							
Element	Total CCZ nodule resource ^a (×10 ⁶ tons)	Global terrestrial reserve base ^b (×10 ⁶ tons)	Global terrestrial resource ^c (×10 ⁶ tons)	Example metal uses			
Manganese	5,992	5,200	ND	Steel, batteries			
Nickel	274	150	ND	Stainless steel, superalloys, wind turbines, batteries			
Copper	226	1,000	5,600	Electrical, electronic, most high-tech products			
Titanium	67	900	1,200	Aerospace, superalloys			
Cobalt	44	13	ND	Batteries, superalloys, electromagnets			
TREE	15.1	128	ND	Turbines, high-tech smartphones etc.			
Molybdenum	12	19	25.4	Steel for strength and hardness			
Vanadium	9.4	38	~63	Steel alloys, jet engines			
Zirconium	6.5	77	ND	Nuclear industry			
Thallium	4.2	0.0007	0.65	Photoresistors, infrared optics			
Lithium	2.8	11	62	Batteries, aircraft			
Yttrium	2.0	0.6	ND	Red phosphor for televisions			
Arsenic	1.4	1.6	ND	Semiconductors			
Tungsten	1.3	6.3	ND	High-strength steel, superalloys, electrodes			
Tellurium	0.08	0.05	ND	Solar cells, superalloys			

CCZ, Clarion–Clipperton Zone; ND, no data; TREE, total rare earth elements. ^aCalculated on the basis of an estimated 21.1 billion metric tons of CCZ nodules in place on the seafloor and the mean chemical composition detailed in TABLE 1. bReserve base is the measured plus indicated resource that includes resources that are currently economic (reserves), marginally economic and some that are currently subeconomic^{129, c}Terrestrial resource is a concentration of a naturally occurring material in or on the Earth's crust that is in such a form and amount that economic extraction is currently or potentially feasible¹³⁰.



Fig. 4 | **The components of a polymetallic-nodule-mining project.** Major components include the collector on the seafloor, the vertical riser system, the surface mining vessel, bulk carriers and the processing plant on land. Multibeam, hydroacoustic mapping with a system mounted on the hull of a vessel provide information about the ocean-floor topography and water depth (bathymetry), and the characteristics of the ocean floor (hard rock, sediments, nodules). Deep-towed systems and autonomous underwater vehicles (AUVs) can get closer to the ocean floor and, thus, provide a higher resolution (smaller footprint). GHG, greenhouse gas; ROV, remotely operated underwater vehicle.

accomplished by one of several methods: pyrometallurgy, which involves smelting the nodules at 1,400–1,500 °C; hydrometallurgy, the chemical dissolution of nodules in sulfuric or hydrochloric acid or in ammonium sulfate and carbonate solutions; and microbiological treatment, such as dissolution by microorganisms⁹⁷.

During pyrometallurgical processing, nodules are smelted in an electric arc furnace under reducing conditions to separate the Mn–Fe–Si slag from a Cu–Ni– Co–Mo metal alloy. The slag can further be treated to produce ferromanganese and silicomanganese for the steel industry and the metal alloy will then undergo hydrometallurgical processing to separate the metals. Finally, the generation of a compact, environmentally stable and toxin-free final slag, applicable to the construction industry, ensures an almost zero-waste processing route for the nodules⁹⁹.

By contrast, many different nodule-processing schemes are solely based on hydrometallurgy. For example, the Cuprion process uses CO as the reducing agent in the presence of ammonium (NH₄) as a lixiviant. Other hydrometallurgical treatments include reduction roasting at 500–750 °C and subsequent leaching with (NH₄)₂CO₃ solution (Caron process), and high-pressure acid leaching or HCl-MgCl₂ leaching (both of which are widely used to process Ni laterites)⁹⁷.

New approaches include the application of microbiological methods based on the use of bacteria or fungi to dissolve and release metals from the oxide lattice^{100,101}, but such methods have, so far, been confined to laboratoryscale experiments. The recovery of trace metals, such as REY and Li, by microbial activity is another area of research, since their extraction could have a positive effect on the economics of nodule processing¹⁰².

Environmental impacts. Before polymetallic-nodule mining enters large-scale operation, it is necessary to consider the possible environmental impacts and how they might be minimized or mitigated. Critically, it is imperative to obtain an adequate understanding of the ecosystems that characterize deep-ocean environments and their interconnectedness. The extraction of marine minerals and the environmental and ecological impacts are a concern in many countries, even in those that are not expected to be directly affected, which could lead to opposition to deep-ocean mining¹⁰³. It is, therefore, important to provide evidence-based risk assessments of deep-ocean mining and to effectively communicate findings to all relevant authorities and countries, as well as the public.

The mining of polymetallic nodules will have biological and geochemical impacts on the ocean floor and overlying water column, the extent and magnitude of which remain uncertain¹⁰⁴. Nodules serve as a hard substrate for a variety of sessile organisms (such as sponges and corals), and octopods have been observed to brood clutches of eggs laid on the stalks of dead sponges attached to nodules¹⁰⁵. The nodule collector is a heavy vehicle that will crush any organisms that are unable to escape and compact the sediment, reducing its habitability for sediment infauna¹⁰⁶⁻¹⁰⁸. Additionally, nodule mining could alter the geochemical composition of the

sediment–water interface and cause a short-term release of potentially toxic metals into the water column¹⁰⁹ (FIG. 4). However, a clear assessment of the potential toxicity of released metals remains a challenge because of the complex composition of polymetallic nodules and sediment particles, as well as the lack of toxicity studies for deep-ocean organisms¹¹⁰. Moreover, the suspension and redeposition of sediment will affect filter-feeding organisms that depend on the supply of nutrient-bearing particles in clean water. The extraction of nodules will, therefore, cause habitat degradation¹¹¹.

The collector systems also pose major problems due to the mobilization of large amounts of sediment. With a mining rate of $0.5 \,\mathrm{m\,s^{-1}}$ and a collector width of $15 \,\mathrm{m}$, 65,000 m3 of sediment is mobilized per day for hydraulic uptake (which impacts the top 10 cm of sediment), resulting in the formation of sediment plumes along the track of the collector (see technical details from REF.⁹¹). The particle plumes might affect not only areas adjacent to the mine sites but also regions at much greater transport distances, depending on prevailing bottom currents and the behaviour of the particles in the dispersing plume. Numerical models published in 2001 indicate that these sediment plumes could have a farreaching impact^{112,113}, although an experimental study published in 2019 demonstrated that aggregation and rapid sinking processes encompass more than 95% of the particles of a sediment plume, thus significantly limiting plume spreading¹¹⁴. However, the effect of the colloidal fraction, which could remain in the water column over much longer timescales and might also transport potentially toxic metals, is not known.

In addition, water and particles from the mining vessel (originating from the dewatering process of the ore on board) will be discharged back into the ocean. By general consensus, such disposal must not take place in surface or shallow waters to avoid interference with biological productivity in the photic zone or redissolution of material in the oxygen minimum zone, but should, instead, be confined to a water depth of at least 1,000 m, preferably close to the ocean floor¹¹⁵. However, no large-scale mining test including the disposal of mine water has taken place as yet. Moreover, the exact composition of the dewatering waste is unclear; predictions of the fate and effects of these discharge waters are also limited¹¹⁶.

As shown by the disturbance and recolonization experiment (DISCOL), and the Joint Programming Initiative Healthy and Productive Seas and Oceans (JPI Oceans mining impact project), nodule mining will have impacts that can be considered permanent on human timescales, if remediation after the removal of nodules (the hard substrate for sessile biota) does not take place^{117,118}. Nodule mining requires large areas to make the enterprise economic11; however, only a certain fraction of each contract area (possibly in the range of 15–20%)^{3,119} will be mined as a first-generation mine site because large parts of the region will have either insufficient nodule coverage or the morphology of the seafloor will preclude harvesting by the mining machine. Hence, organisms are likely to have refugium areas and large areas from which larvae can be dispersed. Nevertheless, even if the overall biomass in an affected area resembles

that of the former undisturbed area, following some period of recovery, the changed habitat may host a different faunal community^{106,108,117}.

Although research on deep-ocean ecosystems^{120,121} has highlighted the high biodiversity and the slow growth and reproduction of deep-ocean fauna, the exact genetic distribution of species (including possible endemic species in certain nodule-bearing areas), as well as the interconnectedness among these populations, are still largely unknown. Furthermore, most studies have primarily focused on benthic fauna, whereas the possible impacts of deep-ocean mining on pelagic and benthopelagic fauna are not often addressed¹²². As a result, it is very difficult to present reliable predictions for the longterm impact of mining on recovery rates and resilience of the deep-ocean ecosystems.

In addition to the marine impact of nodule mining, it is important to consider possible atmospheric and terrestrial environmental consequences to assess the full footprint of deep-ocean mining. For example, greenhouse gas emissions from the mining ship, transport ship and nodule processing on land are all important factors to consider. As long as heavy fuel oil is used for shipping, the release of greenhouse gases and air pollution will be an important issue¹²³. Metallurgical processing of nodules at land-based facilities will also result in the release of greenhouse gases and, possibly, toxic waste¹²⁴. However, the extent of these emissions will depend on the type of technology used and the environmental legislation of the responsible or sponsoring state.

Growth of the deep-ocean-mining industry offers the opportunity to develop green technologies and policies as the industry develops, and concomitantly initiates new strategies to mitigate the environmental effects. Green technologies can be developed for mineral exploration and extraction, as well as ore processing. Development of the deep-ocean-mining industry also provides key opportunities to collect new biological, geological, geochemical and oceanographic data to provide a much clearer understanding of the processes that control the geochemical properties of the ocean floor and water column in mineral-deposit areas.

Finally, deep-ocean mining might help to reduce the pressure on ecologically sensitive ecosystems on land (for example, the deforestation of rainforests to mine laterites for Co and Ni). The replacement of terrestrial mines that impact critical ecosystems with deepocean mines could contribute to fulfilling recent climate treaties and United Nations sustainability goals.

Future perspectives and conclusions

The United States and the European Union, respectively, listed a set of 35 and 27 CRM that are essential to their continued security and prosperity, but are vulnerable to disruption of supply. The sources of CRM required for high-tech, green and alternative energy applications are currently limited and it has become necessary to dig deeper and deeper in terrestrial mines. Deep-ocean mineral deposits, in the form of polymetallic nodules on the seafloor, are rich in many of these CRM. As a result, a complex new industry focused on the recovery of polymetallic nodules from the abyssal plains of the global ocean, and extraction of the metals they contain, is taking shape.

A holistic approach to this developing industry must encompass the concept of 'cradle-to-grave' analysis: assessment of the entire life cycle of product manufacture from exploration and extraction to restoration and/or remediation. This approach is essential and routinely considered as the most constructive way forward for the mining industry. A global strategy for the management of critical metal resources and energy production (such as that implemented by the United Nations) will be required to meet climate and sustainable-development goals^{74,125}. Opportunities exist to develop new green technologies for exploration, mining and ore processing. Application of the precautionary approach, along with adaptive management supported by real-time monitoring, offers the opportunity to ensure the long-term sustainability of this new industry.

The greatest challenge associated with the development of deep-ocean mining is understanding and minimizing its environmental impact, which will determine whether deep-ocean mining should be undertaken and whether it will be of value to societies. In this regard, it is commonly asked whether deep-ocean mining will be more environmentally sustainable than terrestrial mining. However, it is difficult to compare the value of a terrestrial rainforest or grassland with that of a deep-sea ecosystem. What measures should be used? Ecosystem service, biodiversity of the system as a whole, operational CO₂ emissions or respective economic, social and environmental impacts? Certainly, for CO₂ emissions, mining one deep-ocean mineral deposit for three or four metals would have a lower CO₂ footprint than that of three or four terrestrial mines each producing one metal. CO_2 footprint perhaps represents a more easily understood metric than ecosystem-based metrics, but it cannot be considered alone. It is important to note that, although deep-ocean mines will eventually replace some terrestrial mines, especially for critical metals, terrestrial mines will continue to operate well into the future. Both will need to apply best environmental practices using the precautionary approach, which might be easier to achieve in a new industry than to retrofit these concepts to a mature industry that was underway long before the concept of the precautionary approach existed.

Modern societies cannot exist without the materials that are found in polymetallic nodules, and demand for these metals is projected to increase markedly in the future. The question is, however, where should these metals come from? Economic models show that recycling alone is not a viable solution, and that, even in 2050, a substantial part of our economy will depend on primary metal resources^{126,127}. Recycling could become an important part of the answer by 2100, but a much greater focus on education and technology in this arena must take place. The enormous amount of marine mineral resources, and the development of technology to access them, makes deep-ocean contributions to the production of critical minerals seem inevitable. The size of that contribution will, however, depend on other factors, including increased efforts to expand efficient recycling and the success of materials science in finding efficient and high-capacity substitutes for critical materials.

Published online 24 February 2020

 Kuhn, T., Wegorzewski, A., Rühlemann, C. & Vink, A. in *Deep-Sea Mining* (ed. Sharma, R.) 23–63 (Springer, 2017).

An in-depth discussion of the global composition, formation and occurrence of manganese nodules.

- Glasby, G. P. in *Marine Geochemistry* (eds Schulz, H. D. & Zabel M.) 371–427 (Springer, 2006).
 Kuhn, T., Rühlemann, C. & Wiedicke-Hombach, M.
- Kuhn, T., Rühlemann, C. & Wiedicke-Hombach, M. in Marine Minerals: Finding the Right Balance of Sustainable Development and Environmental Protection. Extended Abstracts Vol. 9 (eds Zhou, H. & Morgan, C. L.) (The Underwater Mining Institute, 2012).
- Dymond, J. et al. Ferromanganese nodules from MANOP sites H, S, and R — control of mineralogical and chemical composition by multiple accretionary processes. *Geochim. Cosmochim. Acta* 48, 931–949 (1984).
- 5. Knobloch, A. et al. in *Deep-Sea Mining* (ed. Sharma, R.) 189–212 (Springer, 2017).
- Cronan, D. S. in *Chemical Oceanography* Vol. 5 (eds Riley, J. P. & Chester, R.) 217–263 (Academic, 1976).
 International Seabed Authority. A geological model of
- International seabed Authomy, A geological model of polymetallic nodule deposits in the Clarion-Clipperton Fracture Zone. International Seabed Authority https://www.isa.org.jm/documents/geological-modelpolymetallic-nodule-deposits-clarion-clippertonfracture-zone (2010).
- Hein, J. R., Mizell, K., Koschinsky, A. & Conrad, T. A. Deep-ocean mineral deposits as a source of critical metals for high- and green-technology applications: comparison with land-based resources. *Ore Geol. Rev* 51, 1–14 (2013).
 Details the relationship between critical metals in

deep-ocean mineral deposits to green energy and high-technology applications.

- Cronan, D. S. Deep-sea minerals. *Geoscientist Online* https://www.geolsoc.org.uk/Geoscientist/Archive/ September-2015/Deep-sea-minerals (2015).
- 10. Koschinsky, A. et al. Deep-sea mining: interdisciplinary research on potential environmental, legal, economic,

and societal implications. *Integr. Environ. Assess. Manag.* **14**, 672–691 (2018). **A comprehensive, transdisciplinary discussion of**

the environmental and societal consequences and impacts of future mining of polymetallic nodules, crusts and sulfides.

- Sharma, R. & Smith, S. in *Environmental Issues of Deep-Sea Mining* (ed. Sharma, R.) 3–22 (Springer, 2019).
- Thomson, C. W. & Murray, J. Report on the scientific results of the voyage of H.M.S. Challenger during the years 1872–76 (Printed for Her Majesty's Government, 1895).
- Murray, J. On the distribution of volcanic debris over the floor of the ocean. *Proc. R. Soc. Edinb.* 9, 247–261 (1878).
- von Gümbel, C. W. in *Die am Grunde des Meeres* Vorkommenden Manganknollen 189–209 (Bayerische Akademie der Wissenschaften, 1878).
- Mero, J. L. Ocean-floor manganese nodules. *Economic Geol.* 57, 747–767 (1962).
- 16. Mero, J. L. *The Mineral Resources of the Sea* 1–312 (Elsevier, 1965).
- Buser, W. & Grütter, A. Über die Natur der Manganknollen. Schweiz. Mineral. Petrogr. Mitt. 36, 49–62 (1956).
- Goldberg, E. D. & Arrhenius, G. O. S. Chemistry of Pacific pelagic sediments. *Geochim. Cosmochim. Acta* 13, 153–212 (1958).
- Goldberg, E. D. & Picciotto, E. Thorium determinations in manganese nodules. *Science* 121, 613–614 (1955).
- Riley, J. P. & Sinhaseni, P. Chemical composition of three manganese nodules from the Pacific Ocean. *J. Mar. Res.* **17**, 466–482 (1958).
- Sparenberg, O. A. A historical perspective on deep-sea mining for manganese nodules, 1965–2019. *Extr. Ind. Soc.* 6, 842–854 (2019).
- Morgan, C. L., Nichols, J. A., Selk, B. W., Toth, J. R. & Wallin, C. Preliminary analysis of exploration data from Pacific deposits of manganese nodules. *Mar. Georesour. Geotechnol.* 11, 1–25 (1993).

- National Oceanic and Atmospheric Administration. Deep seabed mining, draft environmental impact statement on issuing exploration licenses to Kennecott Consortium (US Department of Commerce, 1984).
- Sharma, R. in *Deep-Sea Mining* 3–21 (Springer, 2017).
- Bischoff, J. L. & Piper, D. Z. (eds) Marine Geology and Oceanography of the Pacific Manganese Nodule Province (Plenum, 1979).
- von Stackelberg, U. & Marchig, V. Manganese nodule from the equatorial North Pacific Ocean. *Geol. Jahrb.* D87, 123–227 (1987).
- Horn, D. R. (ed.) Ferromanganese Deposits on the Ocean Floor Vol. 293 (National Science Foundation, 1972).
- Glasby, G. P. Marine minerals in the Pacific. Oceanogr. Mar. Biol. Annu. Rev. 24, 11–64 (1986).
- 29. Baturin, G. N. *The Geochemistry of Manganese and Manganese Nodules in the Ocean* (Springer, 1988).
- Usui A. & Moritani T. in Geology and Offshore Mineral Resources of the Central Pacific Basin Vol. 14 (eds. Keating B. H. & Bolton B. R.) 205–223 (Springer, 1992).
- Hein, J. R., Koschinsky, A. & Halliday, A. N. Global occurrence of tellurium-rich ferromanganese crusts and a model for the enrichment of tellurium. *Geochim. Cosmochim. Acta* 67, 1117–1127 (2003).
- Koschinsky, A. & Hein, J. R. Uptake of elements from seawater by ferromanganese crusts: solid-phase associations and seawater speciation. *Mar. Geol.* 198, 331–351 (2003).
- Wegorzewski, A. V. & Kuhn, T. The influence of suboxic diagenesis on the formation of manganese nodules in the Clarion Clipperton nodule belt of the Pacific Ocean. *Mar. Geol.* 357, 123–138 (2014).
- Wegorzewski, A. & Kuhn, T. in Economical, Technological and Environmental Aspects: Cooperative Solutions for Future Deep-Sea Mining (eds Weixler, L. & Goddon, R.) (Underwater Mining Conference, 2017).
- 35. Kashiwabara, T. et al. Chemical processes for the extreme enrichment of tellurium into marine

ferromanganese oxides. *Geochim. Cosmochim. Acta* **131**, 150–163 (2014).

Unravels the surface chemistry and association of tellurium in ferromanganese deposits with both the Fe and Mn phases, the oxidation of the Te on the Mn phase and the process of co-precipitation.

- Koschinsky, A. & Hein, J. R. Marine ferromanganese encrustations: archives of changing oceans. *Elements* 13, 177–182 (2017).
- Jones, D. O. B. et al. Biological responses to disturbance from simulated deep-sea polymetallic nodule mining. *PLOS ONE* 12, e0171750 (2017).
- Heath, G. R. Burial rates, growth rates, and size distributions of deep-sea manganese nodules. *Science* 205, 903–904 (1979).
- Halbach, P., Hebisch, U. & Scherhag, C. Geochemical variations of ferromanganese nodules and crusts from different provinces of the Pacific Ocean and their genetic control. *Chem. Geol.* 34, 3–17 (1981).
- Mewes, K. et al. Diffusive transfer of oxygen from seamount basaltic crust into overlying sediments: an example from the Clarion-Clipperton fracture zone. *Earth Planet. Sci. Lett.* **433**, 215–225 (2016).
- Bau, M. et al. Discriminating between different genetic types of marine ferro-manganese crusts and nodules based on rare earth elements and yttrium. *Chem. Geol.* **381**, 1–9 (2014).
- Bau, M., Koschinsky, A., Dulski, P. & Hein, J. R. Comparison of the partitioning behaviours of yttrium, rare earth elements, and titanium between hydrogenetic marine ferromanganese crusts and seawater. *Geochim. Cosmochim. Acta* 60, 1709–1725 (1996).
- Schmidt, K., Bau, M., Hein, J. R. & Koschinsky, A. Fractionation of the geochemical twins Zr–Hf and Nb–Ta during scavenging from seawater by hydrogenetic ferromanganese crusts. *Geochim. Cosmochim. Acta* 140, 468–487 (2014).
- Banakar, V. K., Pattan, J. N. & Jauhari, P. Size, surface texture, chemical composition and mineralogy interrelations in ferromanganese nodules of central Indian Ocean. *Indian J. Mar. Sci.* 18, 201–203 (1989).
- Kuhn, T., Uhlenkott, K., Vink, A., Rühlemann, C. & Martinez Arbizu, P. in *Seafloor Geomorphology as Benthic Habitat* 2nd edn Ch. 58 (eds Harris, P. & Baker, E.) 933–947 (Elsevier, 2019).

 Halbach, P., Giovanoli, R. & von Borstel, D.
- Halbach, P., Giovanoli, R. & von Borstel, D. Geochemical processes controlling the relationship between Co, Mn, and Fe in early diagenetic deep-sea nodules. *Earth Planet. Sci. Lett.* **60**, 226–236 (1982).
- Halbach, P., Friedrich, G., & von Stackelberg, U. (eds) The Manganese Nodule Belt of the Pacific Ocean: Geological Environment, Nodule Formation and Mining Aspects (Ferdinand Enke, 1988).
 An early, in-depth evaluation of a variety of aspects related to polymetallic nodules and is still considered a benchmark book today.
- Koschinsky, A. & Halbach, P. Sequential leaching of marine ferromanganese precipitates: genetic implications. *Geochim. Cosmochim. Acta* 59, 5113–5132 (1995).
- Usui, A., Mellin, T. Á., Nohara, M. & Yuasa, M. Structural stability of marine 10 Å manganates from the Ogasawara (Bonin) Arc: implication for low-temperature hydrothermal activity. *Mar. Geol.* 86, 41–56 (1989).
- Giovanoli, R. & Arrhenius, G. in *The Manganese* Nodule Belt of the Pacific Ocean: Geological Environment, Nodule Formation and Mining Aspects (eds Halbach, P., Friedrich, G. & von Stackelberg, U.) 20–31 (Ferdinand Enke, 1988).
- Kuhn, T. A., Wegorzewski, C. & Heller, C. R. in Harvesting Seabed Mineral Resources in Harmony with Nature Vol. 6 (eds Morgan, C. L. & Barriga, F. J. A. S.) (The Underwater Mining Institute, 2014).
- Sanderson, B. How bioturbation supports manganese nodules at the sediment-water interface. *Deep Sea Res. A* 32, 1281–1285 (1985).
- Von Stackelberg, U. Significance of benthic organisms for the growth and movement of manganese nodules, Equatorial North Pacific. *Geol. Mar. Lett.* 4, 37–42 (1984).
- Glasby, G. P., Stoffers, P., Sioulas, A., Thijssen, T. & Friedrich, G. Manganese nodule formation in the Pacific Ocean: a general theory. *Geo-Mar. Lett.* 2, 47–53 (1982).
- Nishimura, A. in *Geology and Offshore Mineral* Resources of the Central Pacific Basin Vol. 14 (eds Keating B. H. & Bolton B. R.) 179–203 (Springer, 1992).

- Mewes, K. et al. Impact of depositional and biogeochemical processes on small scale variations in nodule abundance in the Clarion-Clipperton Fracture Zone. Deep Sea Res. Part 191, 125–141 (2014).
- von Stackelberg, U. & Beiersdorf, H. The formation of manganese nodules between the Clarion and Clipperton fracture zones southeast of Hawaii. *Mar. Geol.* **98**, 411–423 (1991).
- Rühlemann, C. & Shipboard Scientific Party. MANGAN 2018: Geology, Biodiversity and Environment of the German License Area for the Exploration of Polymetallic Nodules in the Equatorial NE Pacific. Cruise Report of R/V SONNE cruise SO262 (Bundesanstalt für Geowissenschaften und Rohstoffe, 2019).
- Heller, C., Kuhn, T., Versteegh, G. J. M., Wegorzewski, A. V. & Kasten, S. The geochemical behavior of metals during early diagenetic alteration of buried manganese nodules. *Deep Sea Res. Part I* 142, 16–33 (2018).
- Maeno, M. Y. et al. Sorption behavior of the Pt(II) complex anion on manganese dioxide (δ-MnO₂): a model reaction to elucidate the mechanism by which Pt is concentrated into a marine ferromanganese crust. *Miner. Deposita* 51, 211–218 (2016).
- Takahashi, Y., Ariga, D., Fan, O. & Kashiwabara, T. in Subseafloor Biosphere Linked to Hydrothermal Systems (eds Ishibashi, J., Okino, K. & Sunamura, M.) 39–48 (Springer, 2015).
- Wasylenki, L. E. et al. The molecular mechanism of Mo isotope fractionation during adsorption to birnessite. *Geochim. Cosmochim. Acta* 5, 5019–5031 (2011).
- Manceau, A., Lanson, M. & Takahashi, Y. Mineralogy and crystal chemistry of Mn, Fe, Co, Ni, and Cu in a deep-sea Pacific polymetallic nodule. *Am. Mineral.* 99, 2068–2083 (2014).
- Peacock, C. L. & Moon, E. M. Oxidative scavenging of thallium by birnessite: Explanation for thallium enrichment and stable isotope fractionation in marine ferromanganese precipitates. *Geochim. Cosmochim. Acta* 84, 297–313 (2012).
- Koschinsky, A. et al. Platinum enrichment and phase associations in marine ferromanganese crusts and nodules based on a multi-method approach. *Chem. Geol.* https://doi.org/10.1016/j.chemgeo.2019.119426 (2019).
- (2019).
 Hein, J. R., & Koschinsky, A. in *Treatise on Geochemistry* 2nd edn Vol. 13 Ch. 11 (eds Holland, H. D. & Turekian, K. K.) 273–291 (Elsevier, 2014).
- Kuhn, T., Bau, M., Blum, N. & Halbach, P. Origin of negative Ce anomalies in mixed hydrothermal– hydrogenetic Fe–Mn crusts from the Central Indian Ridge. Earth Planet. Sci. Lett. 163, 207–220 (1998).
- Takahashi, Y., Manceau, A., Geoffroy, N., Matthew, A. M. & Usui, A. Chemical and structural control of the partitioning of Co, Ce, and Pb in marine ferromanganese oxides. *Geochim. Cosmochim. Acta* **71**, 984–1008 (2007).
- Burns, V. M. & Burns, R. G. Authigenic todorokite and phillipsite inside deep-sea manganese nodules. *Am. Mineral.* 63, 827–831 (1978).
- Johnson, E. A. & Post, J. E. Water in the interlayer region of birnessite: importance in cation exchange and structural stability. *Am. Mineral.* **91**, 609–618 (2006).
- Hein, J. R. et al. Critical metals in manganese nodules from the Cook Islands EEZ, abundances and distributions. Ore Geol. Rev. 68, 97–116 (2015).
- Hein, J. R., McIntyre, B. R. & Piper, D. Z. Marine mineral resources of Pacific Islands — a review of the exclusive economic zones of islands of U.S. affiliation, excluding the State of Hawaii. US Geological Survey http://pubs.uses.gov/circ/2005/1286/ (2005).
- Petersen, S. et al. News from the seabed–Geological characteristics and resource potential of deep-sea mineral resources. *Mar. Policy* 70, 175–187 (2016).
- 74. World Bank. Precautionary management of deep sea minerals (World Bank Group, 2017).
- Al Barazi, S., Brandenburg, T., Kuhn, T., Schmidt, M. & Vetter, S. DERA Rohstoffinformationen 36. Kobalt (DERA, 2018).
- Banerji, A. India plans deep dive for seabed minerals. Marine Technology Magazine https:// www.marinetechnologynews.com/news/india-plansseabed-minerals-583447 (2018).
- 77. Tyagi, A., Sudhakar, M. & Chandvale, G. Bulk polymetallic nodule collection in Central Indian Basin — Implications of acoustic survey technologies for site selection and innovations to maximize nodule recovery. *IJOPE* https://www.onepetro.org/conference-paper/ ISOPE-M-09-035 (2009).

- de Moustier, C. Inference of manganese nodule coverage from Sea Beam acoustic backscattering data. *Geophysics* 50, 989–1001 (1985).
- Weydert, M. M. P. Measurements of the acoustic backscatter of selected areas of the deep seafloor and some implications for the assessment of manganese nodule resources. *J. Acoust. Soc. Am.* 88, 350–366 (1990).
- Scanlon, K. M. & Masson, D. G. Fe–Mn nodule field indicated by GLORIA, north of the Puerto Rico Trench. *Geo-Mar. Lett.* 12, 208–213 (1992).
- Masson, D. G. & Scanlon, K. M. Comment on the mapping of iron-manganese nodule fields using reconnaissance sonars such as GLORIA. *Geo-Mar. Lett.* 13, 244–247 (1993).
- Peukert, A. et al. Understanding Mn-nodule distribution and evaluation of related deep-sea mining impacts using AUV-based hydroacoustic and optical data. *Biogeosciences* 15, 2525–2549 (2018).
- Gazis, I.-Z., Schoening, T., Alevizos, E. & Greinert, J. Quantitative mapping and predictive modeling of Mn nodules' distribution from hydroacoustic and optical AUV data linked by random forests machine learning. *Biogeosciences* 15, 7347–7377 (2018).
- Gromoll, L. Geochemical types of Pacific polymetallic nodules: an application of multivariate analysis. Mar. Georesour, Geotechnol. 14, 361–379 (1996).
- Hari, V. N., Kalyan, B., Chitre, M. & Ganesan, V. Spatial modeling of deep-sea ferromanganese nodules with limited data using neural networks. *IEEE J. Ocean. Eng.* 43, 1–18 (2018).
- Kuhn, T., Rühlemann, C. & Knobloch, A. in *Resource* and Environmental Assessments for Seafloor Mining Development (ed. Hong, S.) (Underwater Mining Conference, 2016).
- Lipton, I. T., Nimmo, M. J. & Parianos, J. M. NI 43-101 Technical report. TOML Clarion Clipperton Zone project, Pacific Ocean. *Nautilus Minerals* http:// www.nautilusminerals.com/irm/PDF/1813_0/TOML (2016).
- Boomsma, W. Design of a subsea vehicle propulsion system for soft sediments. Presented at the Underwater Mining Conference (2017).
- Dasselaar, S. Upscaling towards deep-sea mining by 130 meter vertical riser experiments. Presented at the Underwater Mining Conference (2018).
- Hong, S. Technology achievements through 3rd pre-pilot mining test: lifting system. Presented at the Underwater Mining Conference (2016).
- Hong, S. et al. in *Environmental Issues of Deep-Sea Mining* (ed. Sharma, R.) 95–143 (Springer, 2019).
 Xiangyang, L. Progress of Chinese 1000m deep
- Xiangyang, L. Progress of Chinese 1000m deep seabed nodule mining test project. Presented at the Underwater Mining Conference (2019).
- Kuhn, T. et al. Tiefseeförderung von Manganknollen. Schiff & Hafen 5, 78–83 (2011).
- Atmanand, M. A. & Ramadass, G. A. in *Deep-Sea* Mining (ed. Sharma, R.) 305–343 (Springer, 2017).
- van Wijk, J. M. Blue mining: breakthrough solutions for mineral extraction and processing in extreme environments (Blue Mining, 2018).
- environments (Blue Mining, 2018).
 96. Kirchain, R., Roth, R., Field, R. R., III, Muñoz-Royo, C. & Peacock, T. Report to the International Seabed Authority on the development of an economic model and system of payments for the exploitation of polymetallic nodules in the Area (Massachusetts Institute of Technology, 2019).
- portilitate of Technology, 2019).
 97. Das, R. P. & Anand, S. in *Deep-Sea Mining* (ed. Sharma, R.) 365–394 (Springer, 2017).
 A recent overview of the different metallurgical extraction methods applied to Mn nodules.
- Wegorzewski, A., Köpcke, M., Kuhn, T., Sitnikova, M. A. & Wotruba, H. Thermal pre-treatment of polymetallic nodules to create metal (Ni, Cu, Co)-rich individual particles for further processing. *Minerals* 8, 523 (2018).
- Sommerfeld, M., Friedmann, D., Kuhn, T. & Friedrich, B. "Zero-Waste": a sustainable approach on pyrometallurgical processing of manganese nodule slags. *Minerals* 8, 544 (2018).
- Menta, K. D. et al. Bio-dissolution of metals from activated nodules of Indian Ocean. Presented at the International Conference on Frontiers in Mechanochemistry and Mechanical Alloying (2008).
 Mehta, K. D., Das, C. & Pandey, B. B. Leaching of
- Mehta, K. D., Das, C. & Pandey, B. B. Leaching of copper, nickel and cobalt from Indian Ocean manganese nodules by *Aspergillus niger*. *Hydrometallurgy* **105**, 89–95 (2010).
- 102. Mohwinkel, D., Kleint, C. & Koschinsky, A. Phase associations and potential selective extraction methods for selected high-tech metals from ferromanganese

nodules and crusts with siderophores. *Appl. Geochem.* **43**, 13–21 (2014).

- Roche C. & Bice S. Anticipating Social and Community Impacts of Deep Sea Mining 59–80 (Secretariat of the Pacific Community, 2013).
- Cronan, D. S. in *Deep-Sea Polymetallic Nodule* Exploration: Development of Environmental Guidelines 118–154 (International Seabed Authority, 1999).
- 105. Purser, A. et al. Association of deep-sea incirrate octopods with manganese crusts and nodule fields in the Pacific Ocean. *Curr. Biol.* 26, R1268–R1269 (2016).
- Thiel, H. Deep-sea environmental disturbance and recovery potential. *Int. Rev. Gesamten Hydrobiol. Hydrograph.* **77**, 331–339 (1992).
- Borowski, C. & Thiel, H. Deep-sea macrofaunal impacts of a large-scale physical disturbance experiment in the Southeast Pacific. *Deep Sea Res. Part II* 45, 55–81 (1998).
- Jones, D. O. B., Amon, D. J. & Chapman, A. S. A. Mining deep-ocean mineral deposits: What are the ecological risks? *Flements* 14, 325–330 (2018)
- ecological risks? *Elements* 14, 325–330 (2018).
 109. Paul, S. A. L., Gaye, B., Haeckel, M., Kasten, S. & Koschinsky, A. Biogeochemical regeneration of a nodule mining disturbance site: Trace metals, DOC and amino acids in deep-sea sediments and pore waters. *Front. Mar. Sci.* 5, 117 (2018).
- Hauton, C. et al. Identifying toxic impacts of metals potentially released during deep-sea mining — a synthesis of the challenges to quantifying risk. *Front. Mar. Sci.* 4, 368 (2017).
- Niner, H. J. et al. Deep-sea mining with no net loss of biodiversity — an impossible aim. *Front. Mar. Sci.* 5, 53 (2018).
- Jankowski, J. A. & Zielke, W. The mesoscale sediment transport due to technical activities in the deep sea. *Deep Sea Res. Part II* **48**, 3487–3521 (2001).
 Rolinski, S., Segschneider, J. & Sündermann, J.
- 113. Rolinski, S., Segschneider, J. & Sündermann, J. Long-term propagation of tailings from deep-sea mining under variable conditions by means of numerical simulations. *Deep Sea Res. Part II* 48, 3469–3485 (2001).
- 114. Gillard, B. et al. Physical and hydrodynamic properties of deep sea mining-generated, abyssal sediment

plumes in the Clarion Clipperton Fracture Zone (eastern-central Pacific). *Elementa* **7**, 5 (2019).

- Schriever, C. & Thiel, H. Tailings and their disposal in deep-sea mining. *IJOPE* https://www.onepetro.org/ conference-paper/ISOPE-M-13-089 (2013).
- 116. Chung, J. S., Schriever, G., Sharma, R., Yamazaki, T. Deep seabed mining environment: preliminary engineering and environmental assessment. *IJOPE* https://www.onepetro.org/conference-paper/ISOPE-M-01-002 (2001).
- 117. Middelburg, J. et al. (eds) Special issue. Assessing environmental impacts of deep-sea mining-revisiting decade-old benthic disturbances in Pacific nodule areas. *Biogeosciences* https://www.biogeosciences.net/ special_issue942.html (2019).
- Simon-Lledó, E. et al. Biological effects 26 years after simulated deep-sea mining. *Nat. Sci. Rep.* 9, 8040 (2019).
- Volkmann, S. E. & Lehnen, F. Production key figures for planning the mining of manganese nodules. *Mar. Georesour. Geotechnol.* 36, 360–375 (2018).
- 120. Veillette, J. et al. Ferromanganese nodule fauna in the Tropical North Pacific Ocean: Species richness, faunal cover and spatial distribution. *Deep Sea Res. Part I* 54, 1912–1935 (2007).
- 121. De Smet, B. et al. The community structure of deepsea macrofauna associated with polymetallic nodules in the eastern part of the Clarion-Clipperton Fracture Zone. Front. Mar. Sci. 4, 103 (2017).
- 122. Christiansen, B., Denda, A. & Christiansen, S. Potential effects of deep seabed mining on pelagic and benthopelagic biota. *Mar. Policy* https://doi.org/ 10.1016/j.marpol.2019.02.014 (2019).
- 123. Heinrich, L., Markus, T., Koschinsky, A. & Singh, P. Quantifying the fuel consumption, greenhouse gas emissions and air pollution of a potential commercial manganese nodule mining operation. *Mar. Policy* https://doi.org/10.1016/j.marpol.2019.103678 [2019].
- 124. Sen P. K. in *Deep-Sea Mining* (ed. Sharma R.) 395–422 (Springer, 2017).
- 125. Lodge, M. et al. Seabed mining: International Seabed Authority environmental management plan for the Clarion-Clipperton Zone. A partnership approach. *Mar. Policy* **49**, 66–72 (2014).

- McLellan, B. et al. Critical minerals and energy impacts and limitations of moving to unconventional resources. *Resources* 5, 19 (2016).
- 127. Vidal, O., Rostom, F., Francois, C. & Giraud, G. Global trends in metal consumption and supply: The raw material–energy nexus. *Elements* 13, 319–324 (2017).
- Marceau, A., Marcus, M. A. & Grangeon, S. Determination of Mn valence states in mixed-valent manganates by XANES spectroscopy. *Am. Mineral.* 97, 816–827 (2012).
- US Geological Survey. Mineral commodity summaries 2009. US Geological Survey https://www.usgs.gov/ centers/nmic/mineral-commodity-summaries (2009).
- US Geological Survey. Mineral commodity summaries 2019. US Geological Survey https://www.usgs.gov/ centers/nmic/mineral-commodity-summaries (2019).

Acknowledgements

The authors thank S. Whisman for technical help with the preparation of the figures and tables. T.K. is supported by partial funding from the Federal Institute for Geosciences and Natural Resources (BGR) under the grant A-0203002.A.

Author contributions

J.R.H. was responsible for the content, design and compilation of the manuscript. All authors contributed to the preparation and writing of this Review, including the text, figures and tables.

Competing interests

The authors declare no competing interests.

Peer review information

Nature Reviews Earth & Environment thanks R. Sharma, J. González and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

This is a U.S. government work and not under copyright protection in the U.S.; foreign copyright protection may apply 2020